Government of West Bengal **Education Directorate** Bikash Bhawan, Salt Lake, Kolkata - 700 091

Memo No: ED- 827 / 2018

Dated: Kolkata, the 14th September, 2018.

From: The Director of Public Instruction, West Bengal.

To: The Principal/Teacher-in-Charge/Officer-In-Charge of Govt., Govt. aided and erstwhile Sponsored Colleges of West Bengal.

Sub: Preventive measures for fighting Dengue and Chikungunya in the premises of educational institutions as per the Memo No. HIB/M/6-D-08-16/660 Dt.02.03.2017

Ref: Memo No. ED-1391 /2017 dated 23.03.2017, ED-1772/2017 dt.24.5.2018, ED- 1933 / 2017 dt.31.8.2017, ED-1861/ 2017 dated 06.07.2017 and ED-587 / 2018 dated 22.03.2018.

With reference to the earlier letters vide. Memo No. ED-1391 /2017 dated 23.03.2017, ED-1772/2017 dt.24.5.2018, ED- 1933 / 2017 dt.31.8.2017, ED-1861/ 2017 dated 06.07.2017 and ED-587 / 2018 dated 22.03.2018 regarding cleanliness of the premises of educational institutions to prevent Vector Borne diseases, all Principals/Teachers-in-Charge/Officer-in-Charge of Govt., Govt. aided and erstwhile Sponsored colleges are again directed to take necessary measures for prevention & control of Vector-borne disease and general awareness programme among the students regarding preventive measures against Dengue and Chikungunya as per the guidelines set forth by the Directorate of Health Service ,Govt. of West Bengal under the aforesaid memo mentioned above.

To distribute leaflets among the students for awareness generation,

- L To clean the campuses/ premises before commencement of class after a long vacation, П.
- To circulate IEC materials (Provided by Health & F.W. Department) among the III. students.
- To sensitize / engage NSS volunteers in awareness programme regularly. 1V.

This may be treated as extremely urgent. Please send an Action taken report within 15 days from the issuance of this letter to the following e mail: asisghosh.kol@gmail.com

Director of Public Instruction West Bengal

Enclo: Photocopy of Circular as per the Memo No. HIB/M/6-D-08-16/660 Dt.02.03.2017







Government of West Bengal Directorate of Health Services, IBD Branch Swasthya Bhavan, Salt Lake, G. N. -29, Sector- V, Kolkata-700091.

Memo No. HIB/M/6-D-08-16/ 660

Dated - 02 2017

CIRCULAR

From

The Secretary & Mission Director, National Health Mission, Department of H&FW, Govt. of West Bengal

To

The Principal Secretary, Department of Higher Education, Govt. of West Bengal

Subject: Recommended action:

- To maintain the school premises clean & free from breeding source for mosquitoes.
- To increase the general awareness level of students regarding preventive measures against Dengue and Chikungunya.

Sir,

The incidence of Dengue, a mosquito-borne viral disease, has increased in recent years in urban, peri-urban and rural areas. Climatic conditions particularly temperature and rainfall play key roles in the life-cycle, breeding and longevity of dengue vectors and thus in transmission of this deadly disease. The environment in the monsoon season is particularly favorable for dengue mosquito vectors (temperature between 25- 30 Degree Centigrade and relative humidity around 80%), which results in high vector density.

As there is no vaccine or specific drug available against Dengue and Chikungunya infection, vector control', social awareness with ensuring personal protection are significant factors in preventing transmission of both these diseases.

Further, a recent study has revealed that Aedes breeding actually takes place throughout the year. Overhead Tanks and Curing tanks are identified as key containers because they support breeding in both transmission and non-transmission season. During transmission season, vector breeding is spread from these key containers to secondary containers i.e. abandoned containers, coolers, mud pots, blocked drains etc. And thus huge breeding of Aedes mosquitoes sets in.

In this context, may I request you take necessary steps to increase awareness among students/ teachers/ departmental employees, so that the vector control measures against Dengue can be made more intensive.

Following guidelines are enclosed herewith for your kind convenience:

- Annexure I: General Information regarding Dengue & Chikungunya.
- Annexure II: Guideline of Activity for Control & Prevention of Dengue- Chikungunya
- Annexure III: A pictorial instruction of 'Do-s and Don't-s' of Mosquito prevention.
- Annexure IV: IEC Message on Dengue.

I would request you to kindly arrange for appropriate dissemination of the same.

Yours faithfully,

Secretary & Mission Director (NHM) Govt. of West Bengal Dated - 02 3 .2017

Memo No. HIB/M/6-D-08-16/ 660/1 (8)

Copy forwarded for information and necessary action to the:-

- 1) DHS & E.O. Secy, Department of Health & F.W., West Bengal, Swasthya Bhavan, Kol-91.
- Director "NVBDCP, Directorate General of Health Services, GOI, Block-III, Ground Floor, Delhi IT Park, Shastri Park, Delhi-110053.
- 3) Dr.Satyajit Sen; Sr. Regional Director, ROHFW, 27-J.C.Block, Salt Lake, Sector-III, Kolkata-700091.
- 4) Senior P.A. to Additional Chief Secretary, Govt. of West Bengal, Swasthya Bhavan, Kol-91.
- Joint. DHS (PH&CD), West Bengal, Swasthya Bhavan, Kol-91.
- Dy.DHS (Malaria) & SPO, NVBDCP, Swasthya Bhavan, Salt Lake, Kolkata-91.
- Dy. DHS (PH), Swasthya Bhavan, Salt Lake. Kolkata-91.
- 8) Guard File

Jt.DHS(PH&CD) Swasthya Bhavan.

Annexure I

SEMERAL INFORMATION: DENGUE/ DENGUE HAEMORRHAGIC FEVER:

/ECTOR OF DENGUE/DENGUE HAEMORRHAGIC FEVER

- Aedes aegypti is the vector of dengue / dengue haemorrhagic fever.
 - It is a small, black mosquito with white stripes and is approximately 5 mm in size.
- It takes about 7 to 10 days to develop the virus in its body and transmit the disease.

eeding Habit

- Day biter Early morning (8 am- 9 am) and 2 hours before sunset (4 pm- 6 pm)
- Mainly feeds on human beings in domestic and peri-domestic situations
- Bites repeatedly

Resting Habit

- Rests in the domestic and peri-domestic situations
- Rests in the dark corners of houses, on hanging objects like clothes, umbrelia, etc. or under the furniture

Breeding Habits

- Aedes mosquitoes are clean water breeder
- Aedes aegypti mosquito breeds in any type of man-made containers or storage containers having even i small quantity of water
- Eggs of Aedes aegypti can live without water for more than one year

Breeding Places in general

Coolers, Pitch Drums, Water Jars, Mud Pots, Plastic Containers, Buckets, Flower vases, Plant saucers, Over Head Water Tanks, Cisterns, Bottles, Broken Tin Tanks, Deserted Tyres, Roof gutters, Cemetery urns, Bamboo stumps, Coconut shells, Broken egg shells, Bird Pots, Deserted old shoes, Tree holes and many more places where rainwater collects or is stored.

VECTOR CONTROL MEASURES: DO'S AND DON'TS

1. PERSONAL PROPHYLATIC MEASURES

- Use of serosol, mosquito repellent creams, liquids, coils, mats etc to prevent the bites of mosquitoes.
- Remove water from coolers, Air Conditioner Duct/ Water Outlets and other small containers at least once in a week
- Do not wear clothes that expose arms and legs, Wear full sleeve shirts and full pants with socks. Children
 should not be allowed to play in shorts and half sleeved clothes. Try to keep the body covered.
- Use Mosquito nets while sleeping, during day time and night time, to prevent mosquito bite, especially for sleeping infants and young children



COGICAL CONTROL

- Use of larvivorous fishes (like Gambusia, Tilapla, Guppy) in ornamental tanks, fountains, etc.
- Use of blocides

CHEMICAL CONTROL

- Use of chemical larvicides like abate in big breeding containers
- Curing tanks is also peri-domestic containers mainly created by Builders/ Contractors during construction or building and left unattended without demolishing it. Builders/ Contractors should be Responsible for demolishing or mosquito proofing the same by treating these by Temephos or oiling on weekly basis.
- Treat Coolers with Temephos at least twice a week
- Aerosol space spray during day time

ENVIRONMENTAL MANAGEMENT & SOURCE REDUCTION METHODS

- Detection & elimination of mosquito breeding sources on regular basis
- Management of roof tops, porticos and sunshades
- Cover Over Head Tanks (OHTs) properly, by using even domestic cloth
- Proper covering of stored water
- Observation of weekly dry day
- Ensure reliable water supply
- Do not leave solid waste, tyre dumps etc in open. Solid Waste is most crucial breeding habitat during transmission season. So the Administrative Authorities of area should be responsible for the control o same.

HEALTH EDUCATION & COMMUNITY PARTICIPATION

- Sensitize and involve the community for detection of Aedes breeding places and their elimination.
- Impart knowledge to common people regarding the disease and vector through various media sources like T.V., Radio, Cinema slides, Mass Media Campaign with celebrities etc.
- Do Outdoor Publicity: with Hoardings, Posters & Flex Banners in High Risk Blocks
- Distribution of IEC material (like pamphlets, Leaflets, Booklets etc), advertisement in leading News Papers
 wall writings, organizing health camps etc. may also be encouraged.
- Area specific IEC activities may be rigorously promoted by involving key stakeholders, local opinion leaders school students, teachers, PRis etc.



Annexure II Guideline of Activity for Control & Prevention of Dengue-Chikungunya

Activity	Details
Activity Cleanliness Drive at Campus or campus adjacent areas	 Cleanliness Drive in the Campus and adjacent areas for a week It may be treated as a social service/ environment education activity for students. Adoption of neighborhood by institutions for a week to eliminate/ manage breeding sites of mosquitoes. Ensure active participation from adjacent localities Students may distribute hand-written IEC messages/ painted posters etc to the Target Group (TG) This Cleanliness drive should put emphasis on spreading of following IEC messages: Signs and symptoms of Dengue-Chikungunya. Elimination/management of areas where mosquito vector can breed; for e.g. by removal of unused/junk materials like tyres, pots, discarded wet cell batteries, pitch-drums, coconut shells & other scraps from the area, cleaning of stagnant water bodies for channeling of water flow, mud-filling of ditche and other water collections, weakly cleaning of water storages at home covering of over head tanks etc. Protecting oneself from mosquito bites by measures such as regular use of the series o
	 a) Protecting onesen from mosquito ofter of measure of measure of measure of the second of the second

Annexure- III

3

Important Do's and Don'ts



6

Annexure- IV





মনে রাখবে,

জমা জল যেখানে, ভেঙ্গু-বাহক মশার জন্য সেখানে। তাই ব্যক্তির ভিতরে বা আশেপাশে ফেলে রাখা পুরনো টায়ার, ভাঙা ভিম ও ডাবের থোলা, পরিত্যক্ত ব্যাটারির সেল, পিচের ড্রাম, গবাদিপশু

জলের ট্যাঙ্ক, চৌবাচ্চা, এয়ার-কুলার এবং বাড়ির অন্যান্য জলাধারের জল, সপ্তাহে একদিন খালি

ডেব্রুর বাহক এডিস ইজিপ্সাই মশা সাধারণতঃ দিনের বেলায় কামড়ায়। তাই রাতের পাশাপাশি

এমন হলে অবশ্যই ডান্ডনরবাবুর পরামর্শ মতো ওযুষ থাও ও প্রয়োজনে রন্ডপরীক্ষা করাও। সঙ্গে বেশি করে জন্স ও তরলজাতীয় খাবার থেতে হবে। এছাড়াও জুনা হলেই ফেলে না রেখে মত তাড়াতাড়ি সন্থৰ ডাব্ৰুৱবাবুৱ পরামর্শ নাও। সরকারী হাসপাতালে এখন ডেন্সু জ্বরের রোগনির্ণয় ও চিকিৎসার সূযোগ বিনামূলো পাওয়া যায়। মনে রাখবে, অধিকাংশ ডেপ্তু রোগী-ই সময়মতো সাধারণ

স্বাস্থ্য ও পরিবার কল্যাণ দগুর, পশ্চিমবঙ্গ সরকার কর্তৃক প্রচারিত।

ও পোষা পাখির জল খাওয়ার পাত্র বা অন্যান্য অব্যবহৃত পাত্রে জল ভ্রন্মতে দেবেনা।

নর্দমা পরিস্থার রাখে। আেপঝাড় কেট্রে ফেলো। নিয়মিত জঞ্জাল অপসারণ করো। মাঠ বা বাড়ির বাইরে খেলতে গেলে, হাত-পা-ঢাকা হালকা রণ্ডের পোশাক পরো।

IEC Message for Students on Dengue



ডেকু একাট ভাইরাস যটিত স্বর।

মশা-ই ডেন্থ দ্বরের বাহক।

ডেম্বু জ্বরের সাধারণ লক্ষণ হল-

চিকিৎসাই সেরে যায়।

হঠাৎ করে আসা ব্যন্ত, মাথা ব্যাথা চোষের পিছনে, পেশীতে ও গাঁটে ব্যাথা খাব্যরে অরুচি, বমিভাব, পেটে ব্যাথা

বুক-পিঠে-বাহুতে হামের মতো যুসকুড়ি

নাক, মুখ বা মাড়ি থেকে রব্রুক্ষরণ, চামড়ায় কালশিটে

প্রিয় বন্ধুরা, ডেম্বু জ্বর সম্পর্কে সচেতন হও

করে, শুকিয়ে নাও এবং সরসময় ভালোভাবে ঢেকে রাখো। বাড়ির চারপাশে কোনও খানাখন্দে জল জমতে দেবেনা।

শিদের বেলাতেও, ঘুমোনোর সময় অবশ্যই মশারি ব্যবহার করো।